arXiv:1601.01894v1 [math.GR] 8 Jan 2016

On The UNRECOGNIZABILITY BY PRIME GRAPH FOR THE ALMOST SIMPLE GROUP PGL(2, 9)

Ali Mahmoudifar
Department of Mathematics, Tehran North Branch,
Islamic Azad University, Tehran, Iran
e-mail: alimahmoudifar@gmail.com

Abstract

The prime graph of a finite group G is denoted by $\Gamma(G)$. Also G is called recognizable by prime graph if and only if each finite group H with $\Gamma(H) = \Gamma(G)$, is isomorphic to G. In this paper, we classify all finite groups with the same prime graph as $\mathrm{PGL}(2, 9)$. In particular, we present some solvable groups with the same prime graph as $\mathrm{PGL}(2, 9)$.

2000 AMS Subject Classification: 20D05, 20D60, 20D08.
Keywords : Projective general linear group, prime graph, recognition.

1 Introduction

Let n be a natural number. We denote by $\pi(n)$, the set of all prime divisors of n. Also Let G be a finite group. The set $\pi(|G|)$ is denoted by $\pi(G)$. The set of element orders of G is denoted by $\pi_e(G)$. We denote by $\mu(S)$, the maximal numbers of $\pi_e(G)$ under the divisibility relation. The *prime graph* of G is a graph whose vertex set is $\pi(G)$ and two distinct primes p and q are joined by an edge (and we write $p \sim q$), whenever G contains an element of order pq. The prime graph of G is denoted by $\Gamma(G)$. A finite group G is called *recognizable by prime graph* if for every finite group H such that $\Gamma(G) = \Gamma(H)$, then we have $G \cong H$. So G is *recognizable by prime graph* whenever there exists a fin finite group K such that $\Gamma(K) = \Gamma(G)$ in while K is not isomorphic to G.

For the almost simple group $\mathrm{PGL}(2, q)$, there are a lot of results about the recognition by prime graph. In [8], it is proved that if p is a prime number which is not a Mersenne or Fermat prime and $p \neq 11, 19$ and $\Gamma(G) = \Gamma(\mathrm{PGL}(2, p))$, then G has a unique nonabelian composition factor which is isomorphic to $\mathrm{PSL}(2, p)$ and if $p = 13$, then G has a unique nonabelian composition factor which is isomorphic to $\mathrm{PSL}(2, 13)$ or $\mathrm{PSL}(2, 27)$. We know that $\mathrm{PGL}(2, 2^\alpha) \cong \mathrm{PSL}(2, 2^\alpha)$. For the characterization of such simple groups we refer to [9, 10]. In [1], it is proved that if $q = p^\alpha$, where p is an odd prime and α is an odd natural number, then $\mathrm{PGL}(2, q)$ is uniquely determined by its prime graph.

By the above description, we get that the characterization by prime graph of $\mathrm{PGL}(2, p^k)$, where p is an odd prime number and k is even, is an open problem. In this paper as the main result we consider the recognition by prime graph of the almost simple groups $\mathrm{PGL}(2, 3^2)$. Moreover, we construct some solvable group with the same prime graph as $\mathrm{PGL}(2, 3^2)$.

2 Preliminary Results

Lemma 2.1. *Let G be a finite group and $N \trianglelefteq G$ such that G/N is a Frobenius group with kernel F and cyclic complement C. If $(|F|, |N|) = 1$ and F is not contained in $NC_G(N)/N$, then $p|C| \in \pi_e(G)$ for some prime divisor p of $|N|$.*

Lemma 2.2. *Let G be a Frobenius group with kernel F and complement C. Then the following assertions hold:*

(a) F is a nilpotent group.

(b) $|F| \equiv 1 \pmod{|C|}$.

(c) Every subgroup of C of order pq, with p, q (not necessarily distinct) primes, is cyclic. In particular, every Sylow subgroup of C of odd order is cyclic and a Sylow 2-subgroup of C is either cyclic or generalized quaternion group. If C is a non-solvable group, then C has a subgroup of index at most 2 isomorphic to $SL(2,5) \times M$, where M has cyclic Sylow p-subgroups and $(|M|, 30) = 1$.

By using [13, Theorem A] we have the following result:

Lemma 2.3. *Let G be a finite group with $t(G) \geq 2$. Then one of the following holds:*

(a) G is a Frobenius or 2-Frobenius group;

(b) there exists a nonabelian simple group S such that $S \leq \overline{G} := G/N \leq Aut(S)$ for some nilpotent normal π_1-subgroup N of G and $\overline{G}/S$ is a π_1-group.

3 Main Results

Lemma 3.1. *There exists a Frobenius group $G = K : C$, where K is an abelian 3-group and $\pi(C) = \{2, 5\}$, such that $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$.*

Proof. Let F be a finite field with 3^4 elements. Also let V be the additive group of F and H be the multiplicative group $F \setminus \{0\}$. We know that H acts on V by right product. So $G := V \rtimes H$ is a finite group such that $\pi(G) = \{2, 3, 5\}$, since $|V| = 3^4$ and $|H| = 80$. On the other hand H acts fixed point freely on V, so G is a Frobenius group with kernel V and complement H. Since the multiplicative group $F \setminus \{0\}$ is cyclic, H is cyclic too. Therefore, G has an element of order 10, and so the prime graph of G consists just one edge, which is the edge between 2 and 5. This implies that $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$, as desired. □

Lemma 3.2. *There exists a Frobenius group $G = K : C$, where $\pi(K) = \{2, 5\}$ and C is a cyclic 3-group such that $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$.*

Proof. Let F_1 and F_2 be two fields with 2^2 and 5^2 elements, respectively. Let $V := F_1 \times F_2$ be the direct product of the additive groups F_1 and F_2. Also let $H := H_1 \times H_2$, be the direct product of H_1 and H_2, which are the multiplicative groups $F_1 \setminus \{0\}$ and $F_2 \setminus \{0\}$, respectively. We know that H_i acts fixed point freely on F_i, where $1 \leq i \leq 2$. So we define an action of H on V as follows: for each $(h, h') \in H$ and $(g, g') \in V$, we define $(g, g')^{(h,h')} := (hg, h'g')$. It is clear that this definition is well defined. So we may construct a finite group $G = V \rtimes H$. On the other hand H acts fixed point freely on V. So G is a Frobenius group with kernel V and complement H such that $\pi(V) = \{2, 5\}$ and $\pi(H) = \{3\}$. Finally, since V is nilpotent, we get that $\Gamma(G)$ contains an edge between 2 and 5, so $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$. □

Lemma 3.3. *There exists a 2-Frobenius group G with normal series $1 \lhd H \lhd K \lhd G$, such that $\pi(H) = \{5\}$, $\pi(G/K) = \{2\}$ and $\pi(K/H) = \{3\}$, such that $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$.*

Proof. Let F be a field with 5^2 elements and V be its additive group. We know that $F = \{0, 1, \alpha, \alpha^2, \ldots, \alpha^{23}\}$, where α is a generator of the multiplicative group $F \setminus \{0\}$. Hence $|\alpha| = 24$, and so $\beta := \alpha^8$ has order 3. Also $\langle\beta\rangle \cong Z_3$ and $\mathrm{Aut}(Z_3) \cong Z_2$. This argument implies that we may construct a Frobenius group $T := \langle\beta\rangle : \langle\gamma\rangle$, where γ is an involution.

Now we define an action of T on V as follows: for each $\beta^x\gamma^y \in T$ and $v \in V$, $v^{\beta^x\gamma^y} := \beta^x v$, where $1 \leq x \leq 3$ and $1 \leq y \leq 2$. Therefore $G := V : T$ is a 2 Frobenius group with desired properties. □

Theorem 3.4. *Let G be a finite group. Then $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 3^2))$ if and only if G is isomorphic to one of the following groups:*

(1) A Frobenius group $K : C$, where K is an abelian 3-group and $\pi(C) = \{2, 5\}$,

(2) A Frobenius group $K : C$, where $\pi(K) = \{2, 5\}$ and C is a cyclic 3-group,

(3) A 2-Frobenius group with normal series $1 \lhd H \lhd K \lhd G$, such that $\pi(H) \subseteq \{2, 5\}$, $\pi(G/K) = \{2\}$ and $\pi(K/H) = \{3\}$,

(4) Almost simple group $\mathrm{PGL}(2, 3^2)$.

Proof. Throughout the proof, we assume that G is a finite group with the same prime graph as the almost simple group $\mathrm{PGL}(2, 3^2)$. First we note that by [16, Lemma 7], we have:

$$\mu(\mathrm{PGL}(2, 9)) = \{3, 8, 10\}.$$

Hence in $\Gamma(\mathrm{PGL}(2, 3^2))$ (and so in $\Gamma(G)$), there is only one edge which is the edge between 2 and 5 and so 3 is an isolated vertex. This implies that $\Gamma(G)$ has two connected components $\{3\}$ and $\{2, 5\}$. Thus by Lemma 2.3, we get that G is a Frobenius group or 2-Frobenius group or there exists a nonabelian simple group S such that $S \leq G/Fit(G) \leq Aut(S)$. We consider each possibility for G.

Let G be a Frobenius group with kernel K and complement C. We know that K is nilpotent and C is a connected component of the prime graph of G. Also by the above description, 3 is not adjacent to 2 and 5 in $\Gamma(G)$. This shows that either $\pi(K) = \{3\}$ or $\pi(C) = \{3\}$. We consider these cases, separately.

Case 1. Let $\pi(K) = \{3\}$. Hence the order of complement C, is even and so K is an abelain subgroup of G. Also by the above description, $\pi(C) = \{2, 5\}$. Since C is a connected component of $\Gamma(G)$, there is and edge between 2 and 5. So it follows that $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$, which implies groups satisfying in (1).

Case 2. Let $\pi(C) = \{3\}$. Hence $\pi(K) = \{2, 5\}$. Since K is nilpotent, we get that 2 and 5 are adjacent in $\Gamma(G)$. This means $\Gamma(G) = \Gamma(\mathrm{PGL}(2, 9))$ and so we get (2).

Case 3. Let G be a 2-Frobenius group with normal series $1 \lhd H \lhd K \lhd G$. Since $\pi(K/H)$ and $\pi(H) \cup \pi(G/K)$ are the connected components of $\Gamma(G)$, we get that $\pi(K/H) = \{3\}$ and $\pi(H) \cup \pi(G/K) = \{2, 5\}$. This implies (3).

Case 4. Let there exist a nonabelian simple group S, such that $S \leq \bar{G} := G/\mathrm{Fit}(G) \leq \mathrm{Aut}(S)$. Since $\pi(S) \subseteq \pi(G)$, $\pi(S) = \{2, 3, 5\}$. The finite simple groups with this property are classified in [12, Table 8]. So we get that S is isomorphic to one of the simple groups A_5, $\mathrm{PSU}(4, 2)$ and $\mathrm{PSL}(2, 9)(\cong A_6)$.

Subcase 4.1. Let $S \cong A_5$. We know that $\mathrm{Aut}(A_5) = S_5$. So $\bar{G}$ is isomorphic to the alternating group A_5 or the symmetric group S_5. Since in the prime graph of S_5, there is an edge between 2 and 3, hence we get that $\bar{G}$ is not isomorphic to S_5. Thus, $G/\mathrm{Fit(G)} = A_5$. In the prime graph of A_5, 2 and 5 are nonadjacent. Then at least one of the prime numbers 2 or 5, belongs to $\pi(\mathrm{Fit}(G))$.

Let $5 \in \pi(\mathrm{Fit}(G))$. Let F_5 be a Sylow 5-subgroup of $\mathrm{Fit}(G)$. Since F_5 is a characteristic subgroup of $\mathrm{Fit}(G)$ and $\mathrm{Fit}(G)$ is a normal subgroup of G, $F_5 \trianglelefteq G$. On the other hand in alternating group A_5, the subgroup $\langle (12)(34), (13)(24) \rangle : \langle (123) \rangle$ is a Frobenius subgroup isomorphic to $2^2 : 3$. We recall that by the previous argument, $F_5 \trianglelefteq G$, and so G has a subgroup isomorphic to $5^\alpha : (2^2 : 3)$. So by Lemma 2.1, we get that 3 is adjacent to 5, a contradiction.

Subcase 4.2. Let $S \cong \mathrm{PSU}(4, 2)$. By [4], there is an edge between 3 and 2 which is a contradiction.

Subcase 4.3. Let $S \cong \mathrm{PSL}(2, 9)$. Then $\bar{G}$ is isomorphic to $\mathrm{PSL}(2, 9)$ or $\mathrm{PSL}(2, 9) : \langle \theta \rangle$, where θ is a diagonal, field or diagonal-field automorphism of $\mathrm{PSL}(2, 9)$. In particular θ is an involution. If θ is a field or diagonal-field automorphism of $\mathrm{PSL}(2, 9)$, then the semidirect product $\mathrm{PSL}(2, 9) : \langle \theta \rangle$ contains an element of order 6. Hence θ is neither a field automorphism nor a diagonal-field automorphism. Therefore θ is a diagonal automorphism and so $\bar{G} \cong \mathrm{PGL}(2, 9)$.

By the above discussion, $G/\mathrm{Fit}(G) \cong \mathrm{PGL}(2, 9)$. It is enough to prove that $\mathrm{Fit}(G) = 1$. On the contrary, let $r \in \pi(\mathrm{Fit}(G))$. Also let F_r be the Sylow r-subgroup of $\mathrm{Fit}(G)$. Since $\mathrm{Fit}(G)$ is nilpotent, we can write $\mathrm{Fit}(G) = O_{r'}(\mathrm{Fit}(G)) \times F_r$. So if we put $\tilde{G} = G/O_{r'}(\mathrm{Fit}(G))$, then we get that:

$$\mathrm{PGL}(2, 9) \cong \frac{G}{\mathrm{Fit}(G)} \cong \frac{\tilde{G}}{F_r} \cong \frac{\tilde{G}/\Phi(F_r)}{F_r/\Phi(F_r)}.$$

Since $F_r/\Phi(F_r)$ is an elementary abelian group, without loose of generality we may assume that $F := \mathrm{Fit}(G)$ is an elementary abelian r-group and $G/F \cong \mathrm{PGL}(2, 9)$.

If $r = 3$, then by , we conclude that in $\Gamma(G)$, 2 and 3 are adjacent, which is a contradiction. So let $r \neq 3$. Also let S_3 be a Sylow 3-subgroup of $\mathrm{PGL}(2, 9)$. We know that S_3 is not cyclic. On the other hand $F \rtimes S_3$ is a Frobenius group since 3 is an isolated vertex of $\Gamma(G)$. This follows that S_3 is cyclic which is impossible. Therefore $F = 1$ and so $G \cong \mathrm{PGL}(2, 9)$, which completes the proof. □

References

[1] Z. Akhlaghi, M. Khatami and B. Khosravi, Characterization by prime graph of $\mathrm{PGL}(2, p^k)$ where p and k are odd, International Journal of Algebra and Computation 20 (7) (2010) 847-873.

[2] S. Shariati Beynekalae, A. Iranmanesh and M. Foroudi Ghasemabadi, Quasirecognition by prime graph of the simple group, Southeast Asian Bull. Math., $B_n(2)$, 39 (2015) 181–193.

[3] L. Wang, Quasirecognition of $L_{22}(2)$ by Its prime graph, Southeast Asian Bull. Math., 35 (2015) 883–889.

[4] A. A. Buturlakin, Spectra of finite linear and unitary groups, Algebra and Logic 47(2) (2008) 91–99.

[5] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, Atlas of Finite Groups (Oxford University Press, Oxford, 1985).

[6] K. W. Gruenberg and K. W. Roggenkamp, Decomposition of the augmentation ideal and of the relation modules of a finite group, Proc. London Math. Soc. 31(2) (1975) 149–166.

[7] M. Hagie, The prime graph of a sporadic simple group, Comm. Algebra 31(9) (2003) 4405–4424.

[8] M. Khatami, B. Khosravi and Z. Akhlaghi, NCF-distinguishability by prime graph of $PGL(2,p)$, where p is a prime, Rocky Mountain J. Math., to appear.

[9] B. Khosravi, n-Recognition by prime graph of the simple group $PSL(2,q)$, J. Algebra Appl. 7(6) (2008) 735–748.

[10] B. Khosravi, B. Khosravi and B. Khosravi, 2-Recognizability of $PSL(2,p^2)$ by the prime graph, Siberian Math. J. 49(4) (2008) 749.757.

[11] B. Khosravi, B. Khosravi and B. Khosravi, On the prime graph of $PSL(2,p)$ where $p>3$ is a prime number, Acta. Math. Hungarica 116(4) (2007) 295–307.

[12] R. Kogani-Moghadam and A. R. Moghaddamfar, Groups with the same order and degree pattern, *Sci. China Math.*, 55 (4) (2012), 701–720.

[13] A. S. Kondrat'ev, Prime graph components of finite simple groups, Math. USSR-SB. 67(1) (1990) 235–247.

[14] V. D. Mazurov, Characterizations of groups by arithmetic properties, Proceedings of the International Conference on Algebra, Algebra Colloq. 11(1) (2004) 129–140.

[15] V. D. Mazurov, Characterizations of finite groups by sets of their element orders, Algebra Logic 36(1) (1997) 23–32.

[16] A. R. Moghaddamfar and W. J. Shi, The number of finite groups whose element orders is given., Beitrage Algebra Geom., 47(2) (2006) 463–479.

[17] J. S. Williams, Prime graph components of finite groups, J. Algebra 69(2) (1981) 487–513.

[18] A. V. Zavarnitsin, Recognition of finite groups by the prime graph, Algebra Logic 43(4) (2006) 220–231.

[19] K. Zsigmondy, Zur theorie der potenzreste, Monatsh. Math. Phys. 3 (1892) 265–284.